

**Fysioterapeutens undersøgelse ved vestibulær dysfunktion af Kasper Søndergaard. Fysioterapeuten nr. 13 2009.**

REFERENCELISTE

- (1) Hillier SL, Hollohan V. Vestibular rehabilitation for unilateral peripheral vestibular dysfunction. *Cochrane Database Syst Rev* 2007;(4):CD005397.
- (2) Søndergaard K. Vestibulær rehabilitering for kroniske svimmelhed - Et litteraturstudie. 2007.
- (3) Kammerlind AS, Hakansson JK, Skogsberg MC. Effects of balance training in elderly people with nonperipheral vertigo and unsteadiness. *Clin Rehabil* 2001 Oct;15(5):463-70.
- (4) Macias JD, Massingale S, Gerkin RD. Efficacy of vestibular rehabilitation therapy in reducing falls. *Otolaryngol Head Neck Surg* 2005 Sep;133(3):323-5.
- (5) Hall CD, Schubert MC, Herdman SJ. Prediction of fall risk reduction as measured by dynamic gait index in individuals with unilateral vestibular hypofunction. *Otol Neurotol* 2004 Sep;25(5):746-51.
- (6) Whitney SL, Wrisley DM, Marchetti GF, Furman JM. The effect of age on vestibular rehabilitation outcomes. *Laryngoscope* 2002 Oct;112(10):1785-90.
- (7) Basta D, Singbartl F, Todt I, Clarke A, Ernst A. Vestibular rehabilitation by auditory feedback in otolith disorders. *Gait Posture* 2008 Mar 14.
- (8) Malmstrom EM, Karlberg M, Melander A, Magnusson M, Moritz U. Cervicogenic dizziness - musculoskeletal findings before and after treatment and long-term outcome. *Disabil Rehabil* 2007 Aug 15;29(15):1193-205.
- (9) Jacobson GP, Newman CW. The development of the Dizziness Handicap Inventory. *Arch Otolaryngol Head Neck Surg* 1990 Apr;116(4):424-7.
- (10) Gry Hallgren, Louise B. Bockhoff. The Dizziness Handicap Inventory (DHI) Oversættelse til dansk og test af den danske version på ti patienter med svimmelhed. 2008.  
Ref Type: Data File
- (11) Morris AE, Lutman ME, Yardley L. Measuring outcome from vestibular rehabilitation, part II: refinement and validation of a new self-report measure. *Int J Audiol* 2009 Jan;48(1):24-37.
- (12) Smith-Wheelock M, Shepard NT, Telian SA. Physical therapy program for vestibular rehabilitation. *Am J Otol* 1991 May;12(3):218-25.
- (13) Brandt T, Strupp M. General vestibular testing. *Clin Neurophysiol* 2005 Feb;116(2):406-26.
- (14) Halmagyi GM, Weber KP, Aw ST, Todd MJ, Curthoys IS. Impulsive testing of semicircular canal function. *Prog Brain Res* 2008;171:187-94.
- (15) Schubert MC, Tusa RJ, Grine LE, Herdman SJ. Optimizing the sensitivity of the head thrust test for identifying vestibular hypofunction. *Phys Ther* 2004 Feb;84(2):151-8.
- (16) Ramat S, Zee DS, Minor LB. Translational vestibulo-ocular reflex evoked by a "head heave" stimulus. *Ann N Y Acad Sci* 2001 Oct;942:95-113.
- (17) Herdman SJ. *Vestibular Rehabilitation*. F.A. Davis Company; 2007.
- (18) Barber HO. Vestibular neurophysiology. *Otolaryngol Head Neck Surg* 1984 Feb;92(1):55-8.
- (19) Dannenbaum E, Paquet N, Hakim-Zadeh R, Feldman AG. Optimal parameters for the clinical test of dynamic visual acuity in patients with a unilateral vestibular deficit. *J Otolaryngol* 2005 Feb;34(1):13-9.
- (20) Herdman SJ, Tusa RJ, Blatt P, Suzuki A, Venuto PJ, Roberts D. Computerized dynamic visual acuity test in the assessment of vestibular deficits. *Am J Otol* 1998 Nov;19(6):790-6.
- (21) Whitney SL, Marchetti GF, Pritcher M, Furman JM. Gaze stabilization and gait performance in vestibular dysfunction. *Gait Posture* 2009 Feb;29(2):194-8.
- (22) Meretta BM, Whitney SL, Marchetti GF, Sparto PJ,

Muirhead RJ. The five times sit to stand test: responsiveness to change and concurrent validity in adults undergoing vestibular rehabilitation. *J Vestib Res* 2006;16(4-5):233-43.

- (23) Shepard NT, Telian SA. *Practical Management of the Balance Disorders Patient*. 1 ed. 1996.
- (24) Shumway-Cook A, Horak FB. Assessing the influence of sensory interaction of balance. Suggestion from the field. *Phys Ther* 1986 Oct;66(10):1548-50.
- (25) Marchetti GF, Whitney SL, Blatt PJ, Morris LO, Vance JM. Temporal and spatial characteristics of gait during performance of the Dynamic Gait Index in people with and people without balance or vestibular disorders. *Phys Ther* 2008 May;88(5):640-51.
- (26) Kennedy PM, Cressman EK, Carlsen AN, Chua R. Assessing vestibular contributions during changes in gait trajectory. *Neuroreport* 2005 Jul 13;16(10):1097-100.
- (27) Shumway-Cook A, woollacott M. *Motor control: theory and practical applications*. Baltimore: Williams and Wilkins; 1995.
- (28) Whitney SL, Hudak MT, Marchetti GF. The dynamic gait index relates to self-reported fall history in individuals with vestibular dysfunction. *J Vestib Res* 2000;10(2):99-105.
- (29) Whitney SL, Marchetti GF, Schade A, Wrisley DM. The sensitivity and specificity of the Timed "Up & Go" and the Dynamic Gait Index for self-reported falls in persons with vestibular disorders. *J Vestib Res* 2004;14(5):397-409.
- (30) Wrisley DM, Walker ML, Echternach JL, Strasnick B. Reliability of the dynamic gait index in people with vestibular disorders. *Arch Phys Med Rehabil* 2003 Oct;84(10):1528-33.
- (31) Marchetti GF, Whitney SL. Construction and validation of the 4-item dynamic gait index. *Phys Ther* 2006 Dec;86(12):1651-60.
- (32) Wrisley DM, Marchetti GF, Kuharsky DK, Whitney SL. Reliability, internal consistency, and validity of data obtained with the functional gait assessment. *Phys Ther* 2004 Oct;84(10):906-18.
- (33) FUKUDA T. The stepping test: two phases of the labyrinthine reflex. *Acta Otolaryngol* 1959 Mar;50(2):95-108.
- (34) Podsiadlo D, Richardson S. The timed "Up & Go": a test of basic functional mobility for frail elderly persons. *J Am Geriatr Soc* 1991 Feb;39(2):142-8.
- (35) Whitney S, Wrisley D, Furman J. Concurrent validity of the Berg Balance Scale and the Dynamic Gait Index in people with vestibular dysfunction. *Physiother Res Int* 2003;8(4):178-86.
- (36) Dite W, Temple VA. A clinical test of stepping and change of direction to identify multiple falling older adults. *Arch Phys Med Rehabil* 2002 Nov;83(11):1566-71.
- (37) Whitney SL, Marchetti GF, Morris LO, Sparto PJ. The reliability and validity of the Four Square Step Test for people with balance deficits secondary to a vestibular disorder. *Arch Phys Med Rehabil* 2007 Jan;88(1):99-104.