

Cand. scient., ph.d. Ulrik Dalgas. Styrketræning er relevant til patienter med multipel sklerose. Fysioterapeuten nr. 4 2010.

LITTERATUR

1. Kent-Braun JA, Ng AV, Castro M et al. Strength, skeletal muscle composition, and enzyme activity in multiple sclerosis. *J Appl Physiol* 1997; 83: 1998-2004.
2. Ng AV, Kent-Braun JA. Quantitation of lower physical activity in persons with multiple sclerosis. *Med Sci Sports Exerc* 1997; 29: 517-23.
3. Klarlund Pedersen B, Saltin B. Fysisk aktivitet - En håndbog om forebyggelse og behandling. Sundhedsstyrelsen, Center for forebyggelse, 2003.
4. Bronnum-Hansen H, Koch-Henriksen N, Stenager E. Trends in survival and cause of death in Danish patients with multiple sclerosis. *Brain* 2004.
5. Cosman F, Nieves J, Komar L et al. Fracture history and bone loss in patients with MS. *Neurology* 1998; 51: 1161-5.
6. Zorzon M, de Masi R, Nasuelli D et al. Depression and anxiety in multiple sclerosis. A clinical and MRI study in 95 subjects. *J Neurol* 2001; 248: 416-21.
7. Mostert S, Kesselring J. Effects of a short-term exercise training program on aerobic fitness, fatigue, health perception and activity level of subjects with multiple sclerosis. *Mult Scler* 2002; 8: 161-8.
8. Tantucci C, Massucci M, Piperno R et al. Energy cost of exercise in multiple sclerosis patients with low degree of disability. *Mult Scler* 1996; 2: 161-7.
9. Convertino VA, Bloomfield SA, Greenleaf JE. An overview of the issues: physiological effects of bed rest and restricted physical activity. *Med Sci Sports Exerc* 1997; 29: 187-90.
10. Chen WY, Pierson FM, Burnett CN. Force-time measurements of knee muscle functions of subjects with multiple sclerosis. *Phys Ther* 1987; 67: 934-40.
11. de Haan A, de Ruitter CJ, Der Woude LH et al. Contractile properties and fatigue of quadriceps muscles in multiple sclerosis. *Muscle Nerve* 2000; 23: 1534-41.
12. Schwid SR, Thornton CA, Pandya S et al. Quantitative assessment of motor fatigue and strength in MS. *Neurology* 1999; 53: 743-50.
13. Kantarci OH. Genetics and natural history of multiple sclerosis. *Semin Neurol* 2008; 28: 7-16.
14. Kurtzke JF. Epidemiology and etiology of multiple sclerosis. *Phys Med Rehabil Clin N Am* 2005; 16: 327-49.

15. Pilz G, Wipfler P, Ladurner G et al. Modern multiple sclerosis treatment - what is approved, what is on the horizon. *Drug Discov Today* 2008; 13: 1013-25.
16. Liguori M, Marrosu MG, Pugliatti M et al. Age at onset in multiple sclerosis. *Neurol Sci* 2000; 21: S825-S829.
17. Sobocki P, Pugliatti M, Lauer K et al. Estimation of the cost of MS in Europe: extrapolations from a multinational cost study. *Mult Scler* 2007; 13: 1054-64.
18. Ebers GC. Environmental factors and multiple sclerosis. *Lancet Neurol* 2008; 7: 268-77.
19. McDonald WI, Compston A, Edan G et al. Recommended diagnostic criteria for multiple sclerosis: guidelines from the International Panel on the diagnosis of multiple sclerosis. *Ann Neurol* 2001; 50: 121-7.
20. Koch-Henriksen N, Bronnum-Hansen H, Stenager E. Underlying cause of death in Danish patients with multiple sclerosis: results from the Danish Multiple Sclerosis Registry. *J Neurol Neurosurg Psychiatry* 1998; 65: 56-9.
21. Ragonese P, Aridon P, Salemi G et al. Mortality in multiple sclerosis: a review. *Eur J Neurol* 2008; 15: 123-7.
22. Lassmann H, Bruck W, Lucchinetti CF. The immunopathology of multiple sclerosis: an overview. *Brain Pathol* 2007; 17: 210-8.
23. Dalgas U, Stenager E, Ingemann-Hansen T. Multiple sclerosis and physical exercise: recommendations for the application of resistance-, endurance- and combined training. *Mult Scler* 2008; 14: 35-53.
24. DeBolt LS, McCubbin JA. The effects of home-based resistance exercise on balance, power, and mobility in adults with multiple sclerosis. *Arch Phys Med Rehabil* 2004; 85: 290-7.
25. Gutierrez GM, Chow JW, Tillman MD et al. Resistance training improves gait kinematics in persons with multiple sclerosis. *Arch Phys Med Rehabil* 2005; 86: 1824-9.
26. White LJ, McCoy SC, Castellano V et al. Resistance training improves strength and functional capacity in persons with multiple sclerosis. *Mult Scler* 2004; 10: 668-74.
27. Harvey L, Smith A, Jones R. The effect of weighted leg raises on quadriceps strength, EMG parameters and functional activities in people with multiple sclerosis. *Physiotherapy* 1999; 85: 154-61.
28. Kasser S, McCubbin JA. Effects of progressive resistance exercise on muscular strength in adults with multiple sclerosis. *Med Sci Sports Exerc* 1996; 28: S143.
29. Kraft G, Alquist A, Lateur B. Effects of resistive exercise on strength in multiple sclerosis (MS). *Arch Phys Med Rehabil* 1996; 77: 984.
30. Kraft G, Alquist A, Lateur B. Effects of resistive exercise on function in multiple sclerosis (MS). *Arch Phys Med Rehabil* 1996; 77: 984.

31. Fisher NM, Lenox J, Granger CV et al. Effects of an anti-fatiguing exercise program on fatigue and physiological function in patients with Multiple Sclerosis. *Neurology* 2000; 54: A338.
32. Aimet M, Lampichler J, Musil U et al. High and moderate intensities in strength training in multiple sclerosis. *Isokin Exerc Sci* 2006; 14: 153.
33. Ayan PC, Martin S, V, De Souza TF et al. Effects of a resistance training program in multiple sclerosis Spanish patients: a pilot study. *J Sport Rehabil* 2007; 16: 143-53.
34. Dalgas U, Stenager E, Jakobsen J et al. Resistance training improves muscle strength and functional capacity in multiple sclerosis. *Neurology* 2009; 73: 1478-84.
35. de Souza-Teixeira F, Costilla S, Ayan C et al. Effects of resistance training in multiple sclerosis. *Int J Sports Med* 2009; 30: 245-50.
36. Taylor NF, Dodd KJ, Prasad D et al. Progressive resistance exercise for people with multiple sclerosis. *Disabil Rehabil* 2006; 28: 1119-26.
37. Garner DJ, Widrick JJ. Cross-bridge mechanisms of muscle weakness in multiple sclerosis. *Muscle Nerve* 2003; 27: 456-64.
38. Carroll CC, Gallagher PM, Seidle ME et al. Skeletal muscle characteristics of people with multiple sclerosis. *Arch Phys Med Rehabil* 2005; 86: 224-9.
39. Ng AV, Miller RG, Gelinas D et al. Functional relationships of central and peripheral muscle alterations in multiple sclerosis. *Muscle Nerve* 2004; 29: 843-52.
40. Dalgas U. Physical exercise and Multiple Sclerosis - effects of progressive resistance training University of Aarhus, Department of Sport Science, 2009.
41. Staron RS, Malicky ES, Leonardi MJ et al. Muscle hypertrophy and fast fiber type conversions in heavy resistance-trained women. *Eur J Appl Physiol Occup Physiol* 1990; 60: 71-9.
42. Andersen JL, Aagaard P. Myosin heavy chain IIX overshoot in human skeletal muscle. *Muscle Nerve* 2000; 23: 1095-104.
43. Krupp LB, Alvarez LA, LaRocca NG et al. Fatigue in multiple sclerosis. *Arch Neurol* 1988; 45: 435-7.
44. Miller A, Dishon S. Health-related Quality of Life in Multiple Sclerosis: The Impact of Disability, Gender and Employment Status. *Qual Life Res* 2006; 15: 259-71.
45. Smith RM, ey-Steel M, Fulcher G et al. Symptom change with exercise is a temporary phenomenon for people with multiple sclerosis. *Arch Phys Med Rehabil* 2006; 87: 723-7.
46. White AT, Wilson TE, Davis SL et al. Effect of precooling on physical performance in multiple sclerosis. *Mult Scler* 2000; 6: 176-80.

47. Le PC, Bourdoulous S, Beraud E et al. Effect of physical exercise on adoptive experimental auto-immune encephalomyelitis in rats. *Eur J Appl Physiol Occup Physiol* 1996; 73: 130-5.
48. Rossi S, Furlan R, De C, V et al. Exercise attenuates the clinical, synaptic and dendritic abnormalities of experimental autoimmune encephalomyelitis. *Neurobiol Dis* 2009.
49. Berchtold NC, Kessler JP, Pike CJ et al. Estrogen and exercise interact to regulate brain-derived neurotrophic factor mRNA and protein expression in the hippocampus. *Eur J Neurosci* 2001; 14: 1992-2002.
50. Castellano V, White LJ. Serum brain-derived neurotrophic factor response to aerobic exercise in multiple sclerosis. *J Neurol Sci* 2008; 269: 85-91.
51. White LJ, Castellano V, Mc Coy SC. Cytokine responses to resistance training in people with multiple sclerosis. *J Sports Sci* 2006; 24: 911-4.