

# Referencer

## Forsknings-artikel Fysioterapeuten #6, 2024

Af



### MORTEN TANGE KRISTENSEN

Klinisk professor,  
fysioterapeut, ph.d.  
Fysio- og  
Ergoterapiafdelingen  
Bispebjerg og  
Frederiksberg Hospital og  
Institut for  
Klinisk Medicin,  
Københavns Universitet

## Der er evidens for effekten af fysioterapi efter hoftefraktur

- 1 Handoll HHG, Sherrington CPM. **Mobilisation strategies after hip fracture surgery in adults.** Cochrane Database Syst Rev 2004. [doi.org/10.1002/14651858.CD001704.pub3](https://doi.org/10.1002/14651858.CD001704.pub3)
- 2 McDonough CM, Harris-Hayes M, Kristensen MT, et al. **Physical Therapy Management of Older Adults With Hip Fracture.** J Orthop Sports Phys Ther 2021;51. [doi.org/10.2519/jospt.2021.0301](https://doi.org/10.2519/jospt.2021.0301)
- 3 Hulsbæk S, Juhl C, Røpke A, et al. **Exercise Therapy Is Effective at Improving Short- and Long-Term Mobility, Activities of Daily Living, and Balance in Older Patients Following Hip Fracture: A Systematic Review and Meta-Analysis.** Journals Gerontol Ser A 2021;XX:1-11. [doi.org/10.1093/gerona/qlab236](https://doi.org/10.1093/gerona/qlab236)
- 4 Kvalitetsudviklingsprogram RK. **Dansk Tværfagligt Register for Hoftenære Lårbensbrud 2023.**
- 5 Sundhedsstyrelsen. **Tab af muskelfunktion og muskelmasse – sarkopeni og effekten af fysisk træning.** 2023.
- 6 Kristensen MT, Hulsbæk S, Faber LL, et al. **Knee extension strength measures indicating probable sarcopenia is associated with health-related outcomes and a strong predictor of 1-year mortality in patients following hip fracture surgery.** Geriatr 2021;6. [doi.org/10.3390/geriatrics6010008](https://doi.org/10.3390/geriatrics6010008)
- 7 Christensen ER, Clausen A, Petersen TG, et al. **Excess mortality following a first and subsequent osteoporotic fracture: a Danish nationwide register-based cohort study on the mediating effects of comorbidities.** RMD Open 2023;9:1-12. [doi.org/10.1136/rmdopen-2023-003524](https://doi.org/10.1136/rmdopen-2023-003524)
- 8 Overgaard JA, Kallemsø T, Mangione KK, et al. **Six Versus 12 Weeks of Outpatient Physical Therapy Including Progressive Resistance Training in Cognitively Intact Older Adults After Hip Fracture: A Multicenter Randomized Controlled Trial.** Journals Gerontol Ser A 2021;XX:1-8. [doi.org/10.1093/gerona/qlab256](https://doi.org/10.1093/gerona/qlab256)
- 9 Frandsen CF, Stilling M, Glassou EN, et al. **The majority of community-dwelling hip fracture patients return to independent living with minor increase in care needs: a prospective cohort study.** Arch Orthop Trauma Surg 2022. [doi.org/10.1007/s00402-022-04472-8](https://doi.org/10.1007/s00402-022-04472-8)
- 10 Dyer SM, Crotty M, Fairhall N, et al. **A critical review of the long-term disability outcomes following hip fracture.** BMC Geriatr 2016;16. [doi.org/10.1186/s12877-016-0332-0](https://doi.org/10.1186/s12877-016-0332-0)
- 11 Auais MA, Eilayyan O, Mayo NE. **Extended Exercise Rehabilitation After Hip Fracture Improves Patients' Physical Function: A Systematic Review and Meta-Analysis.** Phys Ther 2012;92:1437-51. [doi.org/10.2522/ptj.20110274](https://doi.org/10.2522/ptj.20110274)
- 12 Palm H, Krashennikoff M, Holck K, et al. **A new algorithm for hip fracture surgery.** Acta Orthop 2012;83:26-30. [doi.org/10.3109/17453674.2011.652887](https://doi.org/10.3109/17453674.2011.652887)
- 13 Foss NB. **Hidden blood loss after surgery for hip fracture.** J Bone Jt Surg – Br Vol 2006;88-B:1053-9. [doi.org/10.1302/0301-620X.88B8.17534](https://doi.org/10.1302/0301-620X.88B8.17534)
- 14 Foss NB, Kristensen MT, Jensen PS, et al. **The effects of liberal versus restrictive transfusion thresholds on ambulation after hip fracture surgery.** Transfusion 2009;49:227-34. [doi.org/10.1111/j.1537-2995.2008.01967.x](https://doi.org/10.1111/j.1537-2995.2008.01967.x)

## Referencer.

### Der er evidens for fysioterapi efter hoftefraktur

- 15 Foss NB, Kristensen MT, Kehlet H. **Anaemia impedes functional mobility after hip fracture surgery.** *Age Ageing* 2008;37:173-8. [doi.org/10.1093/ageing/afm161](https://doi.org/10.1093/ageing/afm161)
- 16 Foss NB, Kristensen MT, Palm H, et al. **Postoperative pain after hip fracture is procedure specific.** *Br J Anaesth* 2009;102:111-6. [doi.org/10.1093/bja/aen345](https://doi.org/10.1093/bja/aen345)
- 17 Kristensen MT. **Hip Fracture-Related Pain Strongly Influences Functional Performance of Patients With an Intertrochanteric Fracture Upon Discharge From the Hospital.** *PM R* 2013;5:135-41. [doi.org/10.1016/j.pmrj.2012.10.006](https://doi.org/10.1016/j.pmrj.2012.10.006)
- 18 Kristensen MT, Kehlet H. **Most patients regain prefracture basic mobility after hip fracture surgery in a fast-track programme.** *Dan Med J* 2012;59.
- 19 Kristensen MT, Bandholm T, Bencke J, et al. **Knee-extension strength, postural control and function are related to fracture type and thigh edema in patients with hip fracture.** *Clin Biomech* 2009;24:218-24. [doi.org/10.1016/j.clinbiomech.2008.10.003](https://doi.org/10.1016/j.clinbiomech.2008.10.003)
- 20 Kristensen MT. **Factors influencing performances and indicating risk of falls using the true Timed Up and Go test time of patients with hip fracture upon acute hospital discharge 2020:1-7.** [doi.org/10.1002/pri.1841](https://doi.org/10.1002/pri.1841)
- 21 Kristensen MT, Foss NB, Ekdahl C, et al. **Prefracture functional level evaluated by the New Mobility Score predicts in-hospital outcome after hip fracture surgery.** *Acta Orthop* 2010;81:296-302. [doi.org/10.3109/17453674.2010.487240](https://doi.org/10.3109/17453674.2010.487240)
- 22 Kronborg L, Bandholm T, Palm H, et al. **Feasibility of progressive strength training implemented in the acute ward after hip fracture surgery.** *PLoS One* 2014;9:12-5. [doi.org/10.1371/journal.pone.0093332](https://doi.org/10.1371/journal.pone.0093332)
- 23 Kristensen MT, Keene DJ, Queirós C. **Early Mobilisation and Exercise After Fragility Fracture, 2024, p. 129-46.** [doi.org/10.1007/978-3-031-33484-9\\_8](https://doi.org/10.1007/978-3-031-33484-9_8)
- 24 Dagnelie PC, Willems PC, Jørgensen NR. **Nutritional status as independent prognostic factor of outcome and mortality until five years after hip fracture: a comprehensive prospective study.** *Osteoporos Int* 2024;35:1273-87. [doi.org/10.1007/s00198-024-07088-3](https://doi.org/10.1007/s00198-024-07088-3)
- 25 Schuetze K, Eickhoff A, Rutetzki KS, et al. **Geriatric patients with dementia show increased mortality and lack of functional recovery after hip fracture treated with hemiprosthesis.** *Eur J Trauma Emerg Surg* 2022;48:1827-33. [doi.org/10.1007/s00068-020-01472-4](https://doi.org/10.1007/s00068-020-01472-4)
- 26 Woodcroft-Brown V, Bell J, Pulle CR, et al. **Patient, surgical and hospital factors predicting actual first-day mobilisation after hip fracture surgery: An observational cohort study.** *Australas J Ageing* 2024;1-9. [doi.org/10.1111/ajag.13312](https://doi.org/10.1111/ajag.13312)
- 27 Ariza-Vega P, Kristensen MT, Martín-Martín L, et al. **Predictors of Long-Term Mortality in Older People With Hip Fracture.** *Arch Phys Med Rehabil* 2015;96:1215-21. [doi.org/10.1016/j.apmr.2015.01.023](https://doi.org/10.1016/j.apmr.2015.01.023)

## Referencer.

### Der er evidens for fysioterapi efter hoftefraktur

- 28 Ariza-Vega P, Jiménez-Moleón JJ, Kristensen MT. **Non-weight-bearing status compromises the functional level up to 1 yr after hip fracture surgery.** *Am J Phys Med Rehabil* 2014;93:641-8. [doi.org/10.1097/PHM.0000000000000075](https://doi.org/10.1097/PHM.0000000000000075)
- 29 Viberg B, Kristensen EQ, Gaardsdal T, et al. **A piriformis-preserving posterior approach reduces dislocation rate of the hemiarthroplasty in patients with femoral neck fracture.** *Injury* 2023;54:1727-32. [doi.org/10.1016/j.injury.2023.04.040](https://doi.org/10.1016/j.injury.2023.04.040)
- 30 Münter KH, Clemmesen CG, Foss NB, et al. **Fatigue and pain limit independent mobility and physiotherapy after hip fracture surgery.** *Disabil Rehabil* 2017;0:1-9. [doi.org/10.1080/09638288.2017.1314556](https://doi.org/10.1080/09638288.2017.1314556)
- 31 Zilmer CK, Kristensen MT, Magnusson SP, et al. **Intensified acute in-hospital physiotherapy for patients after hip fracture surgery: a pragmatic, randomized, controlled feasibility trial.** *Disabil Rehabil* 2023;0:1-10. [doi.org/10.1080/09638288.2023.2288672](https://doi.org/10.1080/09638288.2023.2288672)
- 32 Haywood KL, Griffin XL, Achten J, et al. **Developing a core outcome set for hip fracture trials.** *Bone Jt J* 2014;96 B:1016-23. [doi.org/10.1302/0301-620X.96B8.33766](https://doi.org/10.1302/0301-620X.96B8.33766)
- 33 Griffiths F, Mason V, Boardman F, et al. **Evaluating recovery following hip fracture: A qualitative interview study of what is important to patients.** *BMJ Open* 2015;5:1-10. [doi.org/10.1136/bmjopen-2014-005406](https://doi.org/10.1136/bmjopen-2014-005406)
- 34 Hulsbæk, Signe; Røpke, Alice; Damkjær, et al. **Genoptræningsforløbs - beskrivelse for hoftenært brud, Region Hovedstaden.** 2022.
- 35 Kjaer Kristensen P, Thillemann TM, Søballe K, et al. **Are process performance measures associated with clinical outcomes among patients with hip fractures?** A population-based cohort study. *Int J Qual Heal Care* 2016;28:698-708. [doi.org/10.1093/intqhc/mzw093](https://doi.org/10.1093/intqhc/mzw093)
- 36 Kristensen MT, Sheehan KJ, Turabi R. **The relationship between extent of mobilisation within the first postoperative day and 30-day mortality after hip fracture surgery 2024.** [doi.org/10.1177/02692155241231225](https://doi.org/10.1177/02692155241231225)
- 37 Kristensen MT, Öztürk B, Röck ND, et al. **Regaining pre-fracture basic mobility status after hip fracture and association with post-discharge mortality and readmission—a nationwide register study in Denmark.** *Age Ageing* 2019;48:278-84. [doi.org/10.1093/ageing/afy185](https://doi.org/10.1093/ageing/afy185)
- 38 Vesterager JD, Kristensen MT, Pedersen AB. **Loss of pre-fracture basic mobility status at hospital discharge for hip fracture is associated with 30-day post-discharge risk of infections - a four-year nationwide cohort study of 23,309 Danish patients.** *Injury* 2021;52:1833-40. [doi.org/10.1016/j.injury.2021.04.037](https://doi.org/10.1016/j.injury.2021.04.037)
- 39 Perracini MR, Kristensen MT, Cunningham C, et al. **Physiotherapy following fragility fractures.** *Injury* 2018;49:1413-7. [doi.org/10.1016/j.injury.2018.06.026](https://doi.org/10.1016/j.injury.2018.06.026)
- 40 Hagsten B, Svensson O, Gardulf A. **Early individualized postoperative occupational therapy training in 100 patients improves ADL after hip fracture: A randomized trial.** *Acta Orthop Scand* 2004;75:177-83. [doi.org/10.1080/00016470412331294435](https://doi.org/10.1080/00016470412331294435)

## Referencer.

### Der er evidens for fysioterapi efter hoftefraktur

- 41 Hagsten B, Svensson O, Gardulf A. **Health-related quality of life and self-reported ability concerning ADL and IADL after hip fracture: A randomized trial.** Acta Orthop 2006;77:114-9. [doi.org/10.1080/17453670610045786](https://doi.org/10.1080/17453670610045786)
- 42 Lamb SE, Morse RE, Evans JG. **Mobility after proximal femoral fracture: The relevance of leg extensor power, postural sway and other factors.** Age Ageing 1995;24:308-14. [doi.org/10.1093/ageing/24.4.308](https://doi.org/10.1093/ageing/24.4.308)
- 43 Sherrington C, Lord SR, Herbert RD. **A randomized controlled trial of weight-bearing versus non-weight-bearing exercise for improving physical ability after usual care for hip fracture.** Arch Phys Med Rehabil 2004;85:710-6. [doi.org/10.1016/S0003-9993\(03\)00620-8](https://doi.org/10.1016/S0003-9993(03)00620-8)
- 44 Kronborg L, Bandholm T, Palm H, et al. **Effectiveness of acute in-hospital physiotherapy with knee-extension strength training in reducing strength deficits in patients with a hip fracture: A randomised controlled trial.** PLoS One 2017;12:1-18. [doi.org/10.1371/journal.pone.0179867](https://doi.org/10.1371/journal.pone.0179867)
- 45 Sheehan KJ, Goubar A, Almilaji O, et al. **Discharge after hip fracture surgery by mobilisation timing: Secondary analysis of the UK National Hip Fracture Database.** Age Ageing 2021;50:415-22. [doi.org/10.1093/ageing/afaa204](https://doi.org/10.1093/ageing/afaa204)
- 46 Almilaji O, Ayis S, Goubar A, et al. **Frequency, duration, and type of physiotherapy in the week after hip fracture surgery - analysis of implications for discharge home, readmission, survival, and recovery of mobility.** Physiother (United Kingdom) 2023;120:47-59. [doi.org/10.1016/j.physio.2023.03.002](https://doi.org/10.1016/j.physio.2023.03.002)
- 47 Ogawa T, Onuma R, Kristensen MT, et al. **Association between additional weekend rehabilitation and in-hospital mortality in patients with hip fractures.** Bone Jt J 2023;105 E:872-9. [doi.org/10.1302/0301-620X.105B8.BJJ-2022-0890.R3](https://doi.org/10.1302/0301-620X.105B8.BJJ-2022-0890.R3)
- 48 Ogawa T, Onuma R, Sagae H, et al. **Association between additional weekend rehabilitation and functional outcomes in patients with hip fractures: does age affect the effectiveness of weekend rehabilitation?** Eur Geriatr Med 2024. [doi.org/10.1007/s41999-024-00980-5](https://doi.org/10.1007/s41999-024-00980-5)
- 49 Bech RD, Lauritsen J, Ovesen O, et al. **The Verbal Rating Scale Is Reliable for Assessment of Postoperative Pain in Hip Fracture Patients.** Pain Res Treat 2015;2015:1-7. [doi.org/10.1155/2015/676212](https://doi.org/10.1155/2015/676212)
- 50 Mendelsohn ME, Overend TJ, Connelly DM, et al. **Improvement in Aerobic Fitness During Rehabilitation After Hip Fracture.** Arch Phys Med Rehabil 2008;89:609-17. [doi.org/10.1016/j.apmr.2007.09.036](https://doi.org/10.1016/j.apmr.2007.09.036)
- 51 Corna S, Arcolin I, Giardini M, et al. **Addition of aerobic training to conventional rehabilitation after hip fracture: a randomized, controlled, pilot feasibility study.** Clin Rehabil 2021;35:568-77. [doi.org/10.1177/0269215520968694](https://doi.org/10.1177/0269215520968694)
- 52 Cameron ID, Kurrle SE, Sherrington C. **Preventing falls and fall-related injuries in older people.** Med J Aust 2024;1-5. [doi.org/10.5694/mja2.52374](https://doi.org/10.5694/mja2.52374)
- 53 Morin SN, Yan L, Lix LM, et al. **Long-term risk of subsequent major osteoporotic fracture and hip fracture in men and women: a population-based observational study with a 25-year follow-up.** Osteoporos Int 2021;32:2525-32. [doi.org/10.1007/s00198-021-06028-9](https://doi.org/10.1007/s00198-021-06028-9)